

TABLE 2: Dosing Recommendations when Combining Non Nucleoside Reverse Transcriptase Inhibitors and Protease Inhibitors

Protease Inhibitor/Non Nucleoside Reverse Transcriptase Inhibitor Combination	Dosage recommendation
Indinavir with efavirenz or nevirapine	Standard dose for efavirenz and nevirapine. Increase indinavir dosage to 1000mg every eight hours.
Indinavir with delavirdine	Standard dose delavirdine. Decrease indinavir dosage to 600mg every eight hours.
Amprenavir with efavirenz or nevirapine	Standard dose for efavirenz and nevirapine. Increase amprenavir dosage to 1200mg three times daily.
Lopinavir/ritonavir with efavirenz or nevirapine	Standard dose for efavirenz and nevirapine. Increase lopinavir/ritonavir dose to 533mg/133mg (four capsules) twice daily.
Atazanavir with efavirenz (and probably nevirapine, though dosages have not been established with nevirapine)	Standard dose for efavirenz and probably nevirapine. Give atazanavir 300mg once daily AND add ritonavir 100mg once daily
Fosamprenavir with efavirenz (and probably nevirapine, though dosages have not been established with nevirapine)	Standard dose for efavirenz and probably nevirapine. If using fosamprenavir with ritonavir twice daily, give fosamprenavir 700mg with ritonavir 100mg twice daily. If using fosamprenavir with ritonavir once daily, give fosamprenavir 1400mg with ritonavir 300mg once daily.

Data from references 17-21.